

No. DA 23-0572

IN THE SUPREME COURT OF THE STATE OF MONTANA

SCARLET VAN GARDEREN, et al.,

Plaintiffs-Appellees,

v.

STATE OF MONTANA, et al.,

Defendants-Appellants.

On appeal from the Montana Fourth Judicial District Court, Missoula County
Cause No. DV 2023–541, the Honorable Jason Marks, Presiding

**BRIEF OF *AMICI CURIAE* AMERICAN ACADEMY OF PEDIATRICS
AND ADDITIONAL NATIONAL AND STATE MEDICAL AND MENTAL
HEALTH ORGANIZATIONS IN SUPPORT OF APPELLEES**

Jon Moyers (jon@jmoyerslaw.com)
MOYERS KOHN LAW
3936 Ave. B, Suite D
Billings, MT 59102
Phone: (406) 655-4900

D. Jean Veta (jveta@cov.com)*
William R. Isasi (wisasi@cov.com)*
COVINGTON & BURLING, LLP
One CityCenter
850 Tenth St., N.W.
Washington, D.C. 20001
Phone: (202) 662-6000

Counsel for Amici Curiae

**Pro hac vice applications forthcoming*

CORPORATE DISCLOSURE STATEMENT

Pursuant to Montana Rule of Civil Procedure 7.1, the undersigned counsel for the American Academy of Pediatrics (“AAP”), the Academic Pediatric Association, the American Academy of Child & Adolescent Psychiatry (“AACAP”), the American Academy of Family Physicians (“AAFP”), the American Academy of Nursing (“AAN”), the American Association of Physicians for Human Rights, Inc. d/b/a GLMA: Health Professionals Advancing LGBTQ+ Equality (“GLMA”), the American College of Obstetricians and Gynecologists (“ACOG”), the American College of Osteopathic Pediatricians (“ACOP”), the American College of Physicians (“ACP”), the American Medical Association (“AMA”), the American Pediatric Society (“APS”), the Association of American Medical Colleges (“AAMC”), the American Psychiatric Association (“APA”), Association of Medical School Pediatric Department Chairs, Inc. (“AMSPDC”), the Endocrine Society (“ES”), the Montana Chapter of the American Medical Association (“MTAMA”), the Montana Chapter of the American Academy of Pediatrics (“MTAAP”), the National Association of Pediatric Nurse Practitioners (“NAPNAP”), the Pediatric Endocrine Society (“PES”), the Societies for Pediatric Urology (“SPU”), the Society for Adolescent Health and Medicine (“SAHM”), the Society for Pediatric Research (“SPR”), the Society of Pediatric Nurses (“SPN”), and the World Professional Association for Transgender Health (“WPATH”) certify that:

1. AAP, the Academic Pediatric Association, AACAP, AAFP, AAN, GLMA, ACOG, ACOP, ACP, AAMC, AMA, APS, APA, AMSPDC, ES, MTAMA, MTAAP, NAPNAP, PES, SPU, SAHM, SPR, SPN, and WPATH, respectively, have no parent corporation.

2. No corporations hold any stock in AAP, the Academic Pediatric Association, AACAP, AAFP, AAN, GLMA, ACOG, ACOP, ACP, AAMC, AMA, APS, APA, AMSPDC, ES, MTAMA, MTAAP, NAPNAP, PES, SPU, SAHM, SPR, SPN, and WPATH.

TABLE OF CONTENTS

STATEMENT OF INTEREST OF *AMICI CURIAE* 1

SUMMARY OF THE ARGUMENT 3

ARGUMENT 5

I. Understanding Gender Identity and Gender Dysphoria. 6

II. The Widely Accepted Guidelines for Treating Adolescents with Gender Dysphoria Provide for Gender-Affirming Medical Care When Indicated..... 8

 A. The Gender Dysphoria Treatment Guidelines Include Thorough Mental Health Assessments and, for Some Adolescents, Gender-Affirming Medical Care..... 9

 1. The Guidelines Do Not Recommend Gender-Affirming Medical Care for Prepubertal Children..... 10

 2. A Robust Diagnostic Assessment Is Required Before Gender-Affirming Medical Care Is Provided. 10

 3. In Certain Circumstances, the Guidelines Provide for the Use of Gender-Affirming Medical Care to Treat Adolescents with Gender Dysphoria. 11

 B. The Guidelines for Treating Gender Dysphoria Were Developed Through a Robust and Transparent Process, Employing the Same Scientific Rigor That Underpins Other Medical Guidelines..... 15

 C. Scientific Evidence Indicates the Effectiveness of Treating Gender Dysphoria According to the Guidelines. 17

III. The Healthcare Ban Would Irreparably Harm Many Adolescents with Gender Dysphoria by Denying Them the Treatment They Need. 21

CONCLUSION..... 21

CERTIFICATE OF SERVICE 24

TABLE OF AUTHORITIES

	Page(s)
Other Authorities	
Christal Achille et al., <i>Longitudinal Impact of Gender-Affirming Endocrine Intervention on The Mental Health and Wellbeing of Transgender Youths: Preliminary Results</i> , 8 INT’L J PEDIATRIC ENDOCRINOLOGY 1 (2020).....	17, 18
Stewart L. Adelson, <i>Practice Parameter on Gay, Lesbian, or Bisexual Sexual Orientation, Gender Non-Conformity, and Gender Discordance in Children and Adolescents</i> , 51 J. AM. ACAD. CHILD & ADOLESCENT PSYCHIATRY 957 (2020)	7
Zoe Aldridge et al., <i>Long Term Effect of Gender Affirming Hormone Treatment on Depression and Anxiety Symptoms in Transgender People: A Prospective Cohort Study</i> , 9 ANDROLOGY 1808 (2021).....	19
Luke R. Allen et al., <i>Well-Being and Suicidality Among Transgender Youth After Gender-Affirming Hormones</i> , 7 CLINICAL PRAC. PEDIATRIC PSYCH. 302 (2019)	18
AM. PSYCHIATRIC ASS’N, <i>DIAGNOSTIC AND STATISTICAL MANUAL OF MENTAL DISORDERS: DSM-5-TR</i> (2022).....	7
Am. Psychological Ass’n, <i>APA Resolution on Gender Identity Change Efforts</i> , Feb. 2021.....	6
Am. Psychological Ass’n, <i>Guidelines for Psychological Practice with Transgender and Gender Nonconforming People</i> , 70 AMERICAN PSYCHOLOGIST 832 (2015).....	6
Polly Carmichael et al., <i>Short-Term Outcomes of Pubertal Suppression in a Selected Cohort of 12 to 15 Year Old Young People With Persistent Gender Dysphoria in the UK</i> , 16 PLOS ONE e0243894 (2021)	17
Diane Chen et al., <i>Psychosocial Functioning in Transgender Youth after 2 Years of Hormones</i> , 388 NEW ENG. J. MED 240 (2023).....	18, 20, 21

Eli Coleman et al., <i>Standards of Care for the Health of Transgender and Gender Diverse People</i> , 23 <i>Int'l J. Transgender Health</i> S1 (8th ed. 2022).....	<i>passim</i>
F. Comite et al., <i>Short-Term Treatment of Idiopathic Precocious Puberty with a Long-Acting Analogue of Luteinizing Hormone-Releasing Hormone — A Preliminary Report</i> , 305 <i>NEW ENG. J. MED.</i> 1546 (1981).....	13
Rosalia Costa et al., <i>Psychological Support, Puberty Suppression, and Psychosocial Functioning in Adolescents with Gender Dysphoria</i> , 12 <i>J. SEXUAL MED.</i> 2206 (2015).....	17
Annelou L.C. de Vries et al., <i>Puberty Suppression in Adolescents With Gender Identity Disorder: A Prospective Follow-Up Study</i> , 8 <i>J. SEXUAL MED.</i> 2276 (2011).....	17, 20
Annelou L.C. de Vries et al., <i>Young Adult Psychological Outcome After Puberty Suppression And Gender Reassignment</i> , 134 <i>PEDIATRICS</i> 696 (2014).....	17, 18, 20
ENDOCRINE SOC'Y, <i>TRANSGENDER HEALTH: AN ENDOCRINE SOCIETY POSITION STATEMENT</i> (2020).....	8
ENDOCRINE SOC'Y, <i>GUIDELINE METHODOLOGY</i>	16
Amy E. Green et al., <i>Association of Gender-Affirming Hormone Therapy with Depression, Thoughts of Suicide, and Attempted Suicide Among Transgender and Nonbinary Youth</i> , <i>J. ADOLESCENT HEALTH</i> (2021).....	18
Gordon Guyatt et al., <i>GRADE Guidelines: 1. Introduction - GRADE Evidence Profiles and Summary of Findings Tables</i> , 64 <i>J. CLINICAL EPIDEMIOLOGY</i> 383 (2011).....	16
Gordon H. Guyatt et al., <i>GRADE: An Emerging Consensus on Rating Quality of Evidence and Strength of Recommendations</i> , 336 <i>BMJ</i> 924 (2008).....	16
Wylie C. Hembree et al., <i>Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons</i> , 102 <i>J. CLINICAL ENDOCRINOLOGY & METABOLISM</i> 3869 (2017).....	<i>passim</i>

Jody L. Herman et al., <i>Ages of Individuals Who Identify as Transgender in the United States</i> , WILLIAMS INST. (2017).....	8
Michelle M. Johns et al., <i>Transgender Identity and Experiences of Violence Victimization, Substance Use, Suicide Risk, and Sexual Risk Behaviors Among High School Students—19 States and Large Urban School Districts, 2017</i> , U.S. Dep’t of Health and Human Servs., Centers for Disease Control & Prevention, 68 MORBIDITY & MORTALITY WKLY. REP. 67 (2019)	8
Rittakerttu Kaltiala et al., <i>Adolescent Development And Psychosocial Functioning After Starting Cross-Sex Hormones For Gender Dysphoria</i> , 74 NORDIC J. PSYCHIATRY 213 (2020).....	18
Brayden N. Kameg & Donna G. Nativio, <i>Gender Dysphoria In Youth: An Overview For Primary Care Providers</i> , 30 J. AM. ASSOC. NURSE PRAC. 493 (2018).....	7
Laura E. Kuper, et al., <i>Body Dissatisfaction and Mental Health Outcomes of Youth on Gender-Affirming Hormone Therapy</i> , 145 PEDIATRICS e20193006 (2020) 6	17, 18
Diego Lopez de Lara et al., <i>Psychosocial Assessment in Transgender Adolescents</i> , 93 ANALES DE PEDIATRIA 41 (English ed. 2020)	18
James L. Madara, <i>AMA to States: Stop Interfering in Healthcare of Transgender Children</i> , Am. Med. Ass’n (2021)	6
Christy Mallory et al., <i>Conversion Therapy and LGBT Youth</i> , WILLIAMS INST. (2019)	4
Simona Martin et al., <i>Criminalization of Gender-Affirming Care— Interfering with Essential Treatment for Transgender Children and Adolescents</i> , 385 NEW ENG. J. MED. 579 (2021)	5
Anna I.R. van der Miesen, <i>Psychological Functioning in Transgender Adolescents Before and After Gender-Affirmative Care Compared With Cisgender General Population Peers</i> , 66 J. ADOLESCENT HEALTH 699 (2020).....	18

M. Hassan Murad et al., <i>Hormonal Therapy and Sex Reassignment: A Systematic Review and Meta-Analysis of Quality of Life and Psychosocial Outcomes</i> , 72 CLINICAL ENDOCRINOLOGY 214 (2010).....	21
Amit Paley, <i>The Trevor Project 2020 National Survey</i>	8
Ken C. Pang et al., <i>Long-term Puberty Suppression for a Nonbinary Teenager</i> , 145 PEDIATRICS e20191606 (2019).....	14
Jason Rafferty, <i>Ensuring Comprehensive Care and Support for Transgender and Gender-Diverse Children and Adolescents</i> , 142 PEDIATRICS e20182162 (2018).....	4
Stephen M. Rosenthal, <i>Challenges in the Care of Transgender and Gender-Diverse Youth: An Endocrinologist’s View</i> , 17(10) NATURE REV. ENDOCRINOLOGY 581 (2021).....	7, 20
Annemieke S. Staphorsius et al., <i>Puberty Suppression and Executive Functioning: An Fmri-Study in Adolescents with Gender Dysphoria</i> , 6 PSCYHONEUROENDOCRINOLOGY 190 (2015).....	14
Diana M. Tordoff et al., <i>Mental Health Outcomes In Transgender And Nonbinary Youths Receiving Gender-Affirming Care</i> , 5 JAMA NETWORK OPEN e220978 (2022).....	18
Jack L. Turban et al., <i>Access to Gender-Affirming Hormones During Adolescence and Mental Health Outcomes Among Transgender Adults</i> , J. PLOS ONE (2022)	18, 21
Jack L. Turban et al., <i>Pubertal Suppression For Transgender Youth And Risk of Suicidal Ideation</i> , 145 PEDIATRICS e20191725 (2020)	18

STATEMENT OF INTEREST OF *AMICI CURIAE*

Amici curiae are the American Academy of Pediatrics, the Academic Pediatric Association, the American Academy of Child & Adolescent Psychiatry, the Association of American Medical Colleges, the American Academy of Family Physicians, the American Academy of Nursing, the American Association of Physicians for Human Rights, Inc. d/b/a GLMA: Health Professionals Advancing LGBTQ+ Equality, the American College of Obstetricians and Gynecologists, the American College of Osteopathic Pediatricians, the American College of Physicians, the American Medical Association, the American Psychiatric Association, the American Pediatric Society, the Association of Medical School Pediatric Department Chairs, the Endocrine Society, the Montana Chapter of the American Medical Association, the Montana Chapter of the American Academy of Pediatrics, the National Association of Pediatric Nurse Practitioners, the Pediatric Endocrine Society, the Society for Adolescent Health and Medicine, the Society for Pediatric Research, the Society of Pediatric Nurses, the Societies for Pediatric Urology, and the World Professional Association for Transgender Health (collectively, “*amici*”).¹

Amici are professional medical and mental health organizations seeking to

¹ All parties consent to the filing of this brief. *Amici* affirm that no person other than *amici*, their staff, or their counsel made any monetary contributions intended to fund the preparation or submission of this brief.

ensure that all adolescents, including those with gender dysphoria, receive the optimal medical and mental health care they need and deserve. *Amici* represent thousands of healthcare providers who have specific expertise with the issues raised in this brief. The Court should consider *amici*'s brief because it provides important expertise and addresses misstatements about the treatment for transgender adolescents.

SUMMARY OF THE ARGUMENT

On April 17, 2023, Montana Governor Greg Gianforte signed S.B. 99 (the “Healthcare Ban”), enacting a law that effectively bans healthcare providers from providing patients under 18 with critical, medically necessary, evidence-based treatments for gender dysphoria.² The Healthcare Ban makes the provision of such treatments professional misconduct, prohibits the use of public funds for such treatments, and subjects healthcare providers to at least a one year suspension, and private suit.³ Denying such evidence-based medical care to adolescents who meet the requisite medical criteria puts them at risk of significant harm. Below, *amici* provide the Court with an accurate description of the relevant treatment guidelines and summarize the scientific evidence supporting the gender-affirming medical care for adolescents prohibited by the Healthcare Ban.⁴

Gender dysphoria is a clinical condition that is marked by distress due to an incongruence between the patient’s gender identity (i.e., the innate sense of oneself as being a particular gender) and sex assigned at birth. This incongruence can lead

² S.B. 99, 68th Legis. (2023).

³ *Id.* §§ 4-5.

⁴ In this brief, the term “gender-affirming medical care” refers to the use of gonadotropin-releasing hormone (GnRH) analogues and/or hormone therapy to treat gender dysphoria. Because this brief focuses primarily on adolescents, it does not discuss surgeries that are typically available to transgender adults.

to clinically significant distress and impair functioning in many aspects of the patient’s life.⁵ If not treated, or treated improperly, gender dysphoria can result in debilitating anxiety, depression, and self-harm, and is associated with higher rates of suicide. As such, the effective treatment of gender dysphoria saves lives.

The widely accepted recommendation of the medical community, including that of the respected professional organizations participating here as *amici*, is that the well-accepted protocol for treating gender dysphoria is “gender-affirming care.”⁶ Gender-affirming care is care that supports an individual with gender dysphoria as they explore their gender identity—in contrast with efforts to change the individual’s gender identity to match their sex assigned at birth, which are known to be ineffective and harmful.⁷ For adolescents with persistent gender dysphoria that worsens with the onset of puberty, gender-affirming care may include medical care

⁵ See, e.g., Jason Rafferty, *Ensuring Comprehensive Care and Support for Transgender and Gender-Diverse Children and Adolescents*, 142(4) PEDIATRICS e20182162, at 2–3 tbl.1 (2018) [hereinafter, “AAP Policy Statement”], <https://perma.cc/DB5G-PG44>. The American Academy of Pediatrics recently voted to reaffirm the AAP Policy Statement. See Alyson Sulaski Wyckoff, *AAP Reaffirms Gender-Affirming Care Policy, Authorizes Systematic Review of Evidence to Guide Update*, AAP NEWS (Aug. 4, 2023), <https://perma.cc/XS4B-WBLH>. AAP’s review and reaffirmation were undertaken as part of its normal procedures to perform such reviews on a five-year basis.

⁶ AAP Policy Statement, *supra* note 5, at 10.

⁷ See, e.g., Christy Mallory et al., *Conversion Therapy and LGBT Youth*, WILLIAMS INST. (2019), <https://perma.cc/HXY3-UX2J>.

to align their physiology with their gender identity. Empirical evidence indicates that gender-affirming care, including gender-affirming medical care provided to carefully evaluated patients who meet diagnostic criteria, can alleviate clinically significant distress and lead to significant improvements in the mental health and overall well-being of adolescents with gender dysphoria.⁸

The Healthcare Ban disregards this medical evidence by precluding healthcare providers from providing adolescent patients with treatments for gender dysphoria in accordance with the well-accepted protocol. Accordingly, *amici* urge this Court to affirm the district court’s preliminary injunction.

ARGUMENT

This brief first provides background on gender identity and gender dysphoria. It then describes the professionally accepted medical guidelines for treating gender dysphoria as they apply to adolescents, the scientifically rigorous process by which these guidelines were developed, and the evidence that supports the effectiveness of this care for adolescents with gender dysphoria. Finally, the brief explains how the Healthcare Ban would irreparably harm adolescents with gender dysphoria by

⁸ See Simona Martin et al., *Criminalization of Gender-Affirming Care—Interfering with Essential Treatment for Transgender Children and Adolescents*, 385 NEW ENG. J. MED. 579, 580 (2021), <https://perma.cc/BR4F-YLZS> (providing an overview of the scientific basis underlying gender-affirming care and its demonstrated effectiveness in “alleviat[ing] gender dysphoria”).

denying crucial care to those who need it.

I. Understanding Gender Identity and Gender Dysphoria.

A person's gender identity is a person's deep internal sense of belonging to a particular gender.⁹ Most people have a gender identity that aligns with their sex assigned at birth.¹⁰ However, transgender people have a gender identity that does not align with their sex assigned at birth.¹¹ In the United States, it is estimated that approximately 1.4 million individuals are transgender.¹² Of these individuals, approximately 10% are teenagers aged 13 to 17.¹³ Individuals often start to understand their gender identity during prepubertal childhood and adolescence.

Today, there is an increasing understanding that being transgender is a normal variation of human identity.¹⁴ However, many transgender people suffer from gender dysphoria, a serious medical condition in which the patient experiences

⁹ AAP Policy Statement, *supra* note 5, at 2 tbl.1.

¹⁰ See Am. Psychological Ass'n, *Guidelines for Psychological Practice with Transgender and Gender Nonconforming People*, 70 AM. PSYCHOLOGIST 832, 862 (2015), <https://perma.cc/6HR2-9KCM>.

¹¹ See *id.* at 832.

¹² See Jody L. Herman et al., *Ages of Individuals Who Identify as Transgender in the United States*, Williams Inst., Jan. 2017, at 2, <https://perma.cc/C4TA-NR25>.

¹³ See *id.* at 3.

¹⁴ James L. Madara, *AMA to States: Stop Interfering in Healthcare of Transgender Children*, AM. MED. ASS'N (Apr. 26, 2021), <https://perma.cc/BKS6-QFQ8>; see also Am. Psychological Ass'n, *APA Resolution on Gender Identity Change Efforts*, Feb. 2021, at 4, <https://perma.cc/M22K-PBUZ>.

significant distress that can lead to “impairment in peer and/or family relationships, school performance, or other aspects of their life.”¹⁵ Gender dysphoria is a formal diagnosis under the American Psychiatric Association’s Diagnostic and Statistical Manual (DSM-5-TR).¹⁶

Adolescents with gender dysphoria are not expected to identify later as their sex assigned at birth.¹⁷ Instead, “[l]ongitudinal studies have indicated that the emergence or worsening of gender dysphoria with pubertal onset is associated with a very high likelihood of being a transgender adult.”¹⁸

If untreated or inadequately treated, gender dysphoria can cause depression, anxiety, self-harm, and suicidality.¹⁹ Indeed, over 60% of transgender adolescents

¹⁵ AAP Policy Statement, *supra* note 5, at 3.

¹⁶ See AM. PSYCHIATRIC ASS’N, DIAGNOSTIC AND STATISTICAL MANUAL OF MENTAL DISORDERS: DSM-5-TR AT 512–13 (2022).

¹⁷ See, e.g., Stewart L. Adelson, *Practice Parameter on Gay, Lesbian, or Bisexual Sexual Orientation, Gender Nonconformity, and Gender Discordance in Children and Adolescents*, 51 J. AM. ACAD. CHILD & ADOLESCENT PSYCHIATRY 957, 964 (2020), <https://pubmed.ncbi.nlm.nih.gov/22917211> (“In contrast, when gender variance with the desire to be the other sex is present in adolescence, this desire usually does persist through adulthood”).

¹⁸ Stephen M. Rosenthal, *Challenges in the Care of Transgender and Gender-Diverse Youth: An Endocrinologist’s View*, 17(10) NATURE REV. ENDOCRINOLOGY 581, 585 (Oct. 2021), <https://pubmed.ncbi.nlm.nih.gov/34376826>.

¹⁹ See Brayden N. Kameg & Donna G. Nativio, *Gender Dysphoria In Youth: An Overview For Primary Care Providers*, 30 J. AM. ASSOC. NURSE PRAC. 493 (2018), <https://pubmed.ncbi.nlm.nih.gov/30095668>.

and young adults reported having engaged in self-harm during the preceding 12 months, and over 75% reported symptoms of generalized anxiety disorder in the preceding two weeks.²⁰ Even more troubling, more than 50% of this population reported having seriously considered attempting suicide,²¹ and more than one in three transgender adolescents reported having attempted suicide in the preceding 12 months.²²

II. The Widely Accepted Guidelines for Treating Adolescents with Gender Dysphoria Provide for Gender-Affirming Medical Care When Indicated.

The widely accepted view of the professional medical community is that gender-affirming care is the appropriate treatment for gender dysphoria and that, for some adolescents, gender-affirming medical care is necessary.²³ This care greatly reduces the negative physical and mental health consequences that result when

²⁰ See Amit Paley, *The Trevor Project 2020 National Survey*, at 1, <https://perma.cc/JB6T-49XF>.

²¹ See *id.* at 2.

²² See Michelle M. Johns et al., Centers for Disease Control & Prevention, U.S. Dep't of Health and Human Servs., *Transgender Identity and Experiences of Violence Victimization, Substance Use, Suicide Risk, and Sexual Risk Behaviors Among High School Students—19 States and Large Urban School Districts, 2017*, 68 MORBIDITY & MORTALITY WKLY. REP. 67, 70 (2019), <https://perma.cc/7ZKM-F4SS>.

²³ See, e.g., ENDOCRINE SOC'Y, *TRANSGENER HEALTH: AN ENDOCRINE SOCIETY POSITION STATEMENT* (2020), <https://perma.cc/7L4P-VWME>.

gender dysphoria is untreated.²⁴

A. The Gender Dysphoria Treatment Guidelines Include Thorough Mental Health Assessments and, for Some Adolescents, Gender-Affirming Medical Care.

The treatment protocols for gender dysphoria are laid out in established, evidence-based clinical guidelines: (i) the Endocrine Society Clinical Practice Guideline for Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons, and (ii) the WPATH Standards of Care for the Health of Transgender and Gender-Diverse People (together, the “Guidelines”).²⁵ The Guidelines have been developed by expert clinicians and researchers who have worked with patients with gender dysphoria for many years.

The Guidelines provide that all youth with gender dysphoria should be evaluated, diagnosed, and treated by a qualified health care professional (“HCP”). Further, the Guidelines provide that each patient who receives gender-affirming care should receive only evidence-based, medically necessary, and appropriate care that is tailored to the patient’s individual needs.

²⁴ *See id.*

²⁵ Wylie C. Hembree et al., *Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons*, 102 J. CLINICAL ENDOCRINOLOGY & METABOLISM 3869 (2017) [hereinafter, “Endocrine Soc’y Guidelines”], <https://perma.cc/34KY-2LDF>; Eli Coleman et al., *Standards of Care for the Health of Transgender and Gender Diverse People* 23 INT’L J. TRANSGENDER HEALTH S1 (8th ed. 2022) [hereinafter, “WPATH Guidelines”], <https://perma.cc/7SU3-RPK9>.

1. The Guidelines Do Not Recommend Gender-Affirming Medical Care for Prepubertal Children.

For prepubertal children with gender dysphoria, the Guidelines provide for mental health care and support for the child and their family, such as through psychotherapy and social transitioning.²⁶ The Guidelines do *not* recommend that prepubertal children with gender dysphoria receive gender-affirming medical care or surgeries.²⁷

2. A Robust Diagnostic Assessment Is Required Before Gender-Affirming Medical Care Is Provided.

In contrast to prepubertal children, the Guidelines do contemplate the possibility that transgender adolescents with gender dysphoria could receive gender-affirming medical care, provided certain criteria are met. According to the Guidelines, gender-affirming medical care should be provided only after a thorough evaluation by a HCP who: (1) is licensed by their statutory body and holds a master's degree or equivalent in a relevant clinical field; (2) has expertise and received theoretical and evidence-based training in child, adolescent, and family mental health; (3) has expertise and received training in gender identity

²⁶ See WPATH Guidelines, *supra* note 25, at S73–S74; Endocrine Soc’y Guidelines, *supra* note 25, at 3877–78.

²⁷ See WPATH Guidelines, *supra* note 25, at S64, S67; Endocrine Soc’y Guidelines, *supra* note 25, at 3871.

development, gender diversity in children and adolescents, can assess capacity to consent, and possesses knowledge about gender diversity across the life span; (4) has expertise and received training in autism spectrum disorders and other neurodevelopmental presentations, or collaborates with a developmental disability expert when working with neurodivergent patients; and (5) continues engagement in professional development in areas relevant to gender diverse children, adolescents, and families.²⁸

Prior to developing a treatment plan, the HCP should conduct a “comprehensive biopsychosocial assessment” of the adolescent patient.²⁹ The HCP conducts this assessment to “understand the adolescent’s strengths, vulnerabilities, diagnostic profile, and unique needs,” so that the resulting treatment plan is appropriately individualized.³⁰ This assessment must be conducted collaboratively with the patient and their caregiver(s).³¹

3. In Certain Circumstances, the Guidelines Provide for the Use of Gender-Affirming Medical Care to Treat Adolescents with Gender Dysphoria.

For youths with gender dysphoria that continues into adolescence—after the

²⁸ See WPATH Guidelines, *supra* note 25, at S49.

²⁹ *Id.* at S50.

³⁰ *Id.*

³¹ *Id.*

onset of puberty—the Guidelines provide that, in addition to mental health care, gender-affirming medical care may be indicated. Before an adolescent may receive any gender-affirming medical care for treating gender dysphoria, the Guidelines collectively provide that a qualified HCP must determine that: (1) the adolescent meets the diagnostic criteria of gender dysphoria or gender incongruence according to an established taxonomy;³² (2) the adolescent has demonstrated a sustained and persistent pattern of gender nonconformity or gender dysphoria; (3) the adolescent has demonstrated the emotional and cognitive maturity required to provide informed consent for treatment; (4) any coexisting psychological, medical, or social problems that could interfere with diagnosis, treatment, or the adolescent’s ability to consent have been addressed; (5) the adolescent has been informed of the reproductive effects of treatment in the context of their stage in pubertal development and discussed fertility preservation options; and (6) the adolescent has reached Tanner stage 2 of puberty to initiate pubertal suppression.³³ Further, a pediatric endocrinologist or other clinician experienced in pubertal assessment must (7) agree with the indication for treatment, (8) confirm the patient has started puberty, and (9)

³² Endocrine Soc’y Guidelines, *supra* note 25, at 3876; WPATH Guidelines, *supra* note 25, at S47, S48.

³³ WPATH Guidelines, *supra* note 25, at S59–65.

confirm that there are no medical contraindications.³⁴

If all of the above criteria are met, and the patient and their parents provide informed consent, gonadotropin-releasing hormone (GnRH) analogues, or “puberty blockers,” may be offered beginning at the onset of puberty.³⁵ The purpose of puberty blockers is to delay pubertal development until adolescents are old enough and have had sufficient time to make more informed decisions about whether to pursue further treatments.³⁶ Puberty blockers also can make pursuing transition later in life easier, because they prevent irreversible bodily changes such as protrusion of the Adam’s apple or breast growth.³⁷ Puberty blockers have well-known efficacy and side-effect profiles.³⁸ Their effects are generally reversible, and when a patient discontinues their use, the patient resumes endogenous puberty.³⁹ In fact, puberty blockers have been used by pediatric endocrinologists for more than 40 years for the treatment of precocious puberty.⁴⁰ The risks of any serious adverse effects from

³⁴ Endocrine Soc’y Guidelines, *supra* note 25, at 3878 tbl.5.

³⁵ WPATH Guidelines, *supra* note 25, at S64; Martin, *supra* note 8, at 2.

³⁶ WPATH Guidelines, *supra* note 25, at S112.

³⁷ See AAP Policy Statement, *supra* note 5, at 5.

³⁸ See Martin, *supra* note 8, at 2.

³⁹ See *id.*

⁴⁰ See F. Comite et al., *Short-Term Treatment of Idiopathic Precocious Puberty with a Long-Acting Analogue of Luteinizing Hormone-Releasing Hormone — A Preliminary Report*, 305 NEW ENG. J. MED. 1546 (1981).

puberty blockers are exceedingly rare when provided under clinical supervision.⁴¹

Later in adolescence—and if the criteria below are met—hormone therapy may be used to initiate puberty consistent with the patient’s gender identity.⁴² Hormone therapy involves using gender-affirming hormones to allow adolescents to develop secondary sex characteristics consistent with their gender identity.⁴³ Hormone therapy is only prescribed when a qualified mental health professional has confirmed the persistence of the patient’s gender dysphoria, the patient’s mental capacity to consent to the treatment, and that any coexisting problems have been addressed.⁴⁴ A pediatric endocrinologist or other clinician experienced in pubertal induction must also agree with the indication, and the patient and their parents or guardians must be informed of the potential effects and side effects and give their informed consent.⁴⁵ Although some of the changes caused by hormone therapy

⁴¹ See, e.g., Annemieke S. Staphorsius et al., *Puberty Suppression and Executive Functioning: An Fmri-Study in Adolescents with Gender Dysphoria*, 6 *PSYCHONEUROENDOCRINOLOGY* 190 (2015), <https://pubmed.ncbi.nlm.nih.gov/25837854> (no adverse impact on executive functioning); Ken C. Pang et al., *Long-term Puberty Suppression for a Nonbinary Teenager*, 145(2) *PEDIATRICS* e20191606 (2019), <https://perma.cc/VP47-UA9M> (exceedingly low risk of delayed bone mineralization from hormone treatment).

⁴² Martin, *supra* note 8, at 2.

⁴³ See AAP Policy Statement, *supra* note 5, at 6.

⁴⁴ Endocrine Soc’y Guidelines, *supra* note 25, at 3878 tbl.5.

⁴⁵ See *id.*

become irreversible after those secondary sex characteristics are fully developed, others are partially reversible if the patient discontinues use of the hormones.⁴⁶

The Guidelines contemplate that the prescription of puberty blockers and/or hormone therapy be coupled with education on the safe use of such medications and close monitoring to mitigate any potential risks.⁴⁷ Decisions regarding the appropriate treatment for each patient with gender dysphoria are made in consultation with the patient, their parents or guardians, and the medical and mental health care team. There is “no one-size-fits-all approach to this kind of care.”⁴⁸

B. The Guidelines for Treating Gender Dysphoria Were Developed Through a Robust and Transparent Process, Employing the Same Scientific Rigor That Underpins Other Medical Guidelines.

The Guidelines are the product of careful and robust deliberation following the same types of processes—and subject to the same types of rigorous requirements—as other guidelines promulgated by *amici* and other medical organizations.

For example, the Endocrine Society’s Guidelines were developed following a 26-step, 26-month drafting, comment, and review process.⁴⁹ The Endocrine Society

⁴⁶ See AAP Policy Statement, *supra* note 5, at 5–6.

⁴⁷ See Endocrine Soc’y Guidelines, *supra* note 25, at 3871, 3876.

⁴⁸ Martin, *supra* note 8, at 1.

⁴⁹ See, e.g., Endocrine Soc’y Guidelines, *supra* note 25, at 3872–73 (high-level overview of methodology).

imposes strict evidentiary requirements based on the internationally recognized Grading of Recommendations Assessment, Development and Evaluation (GRADE) system.⁵⁰ That GRADE assessment is then reviewed, re-reviewed, and reviewed again by multiple, independent groups of professionals.⁵¹ Reviewers are subject to strict conflict of interest rules, and there is ample opportunity for feedback and debate through the years-long review process.⁵² Further, the Endocrine Society continually reviews its own guidelines and recently determined that the 2017 transgender care guidelines continue to reflect the best, most up-to-date available evidence.

First published in 1979, the WPATH Standards of Care are currently in their 8th Edition. The current Standards of Care are the result of a robust drafting, comment, and review process that collectively took five years.⁵³ The draft guidelines went through rigorous review and were publicly available for discussion

⁵⁰ See Gordon Guyatt et al., *GRADE Guidelines: 1. Introduction - GRADE Evidence Profiles and Summary of Findings Tables*, 64 J. CLINICAL EPIDEMIOLOGY 383 (2011), <https://perma.cc/66FA-6MT6>; Gordon H. Guyatt et al., *GRADE: An Emerging Consensus on Rating Quality of Evidence and Strength of Recommendations*, 336 BMJ 924 (2008), <https://perma.cc/4J7F-3Z62>.

⁵¹ ENDOCRINE SOC'Y, GUIDELINE METHODOLOGY, <https://perma.cc/9NK4-HNNX>.

⁵² See *id.*

⁵³ See WPATH Guidelines, *supra* note 25, at S247-51.

and debate, receiving a total of 2,688 comments.⁵⁴ 119 authors were ultimately involved in the final draft, including feedback from experts in the field as well as from transgender individuals and their families.⁵⁵

C. Scientific Evidence Indicates the Effectiveness of Treating Gender Dysphoria According to the Guidelines.

Multiple studies indicate that adolescents with gender dysphoria who receive gender-affirming medical care experience improvements in their overall well-being.⁵⁶ A number of studies have been published that investigated the use of puberty blockers on adolescents with gender dysphoria,⁵⁷ and/or the use of hormone

⁵⁴ *See id.*

⁵⁵ *See id.*

⁵⁶ *See* Martin, *supra* note 8, at 2.

⁵⁷ *See, e.g.,* Christal Achille et al., *Longitudinal Impact of Gender-Affirming Endocrine Intervention on the Mental Health and Wellbeing of Transgender Youths: Preliminary Results*, 8 INT’L J PEDIATRIC ENDOCRINOLOGY 1–5 (2020), <https://perma.cc/K5SR-EE3G>; Polly Carmichael et al., *Short-Term Outcomes of Pubertal Suppression in a Selected Cohort of 12 to 15 Year Old Young People with Persistent Gender Dysphoria in the UK*, 16 PLOS ONE e0243894 (2021), <https://doi.org/10.1371/journal.pone.0243894>; Rosalia Costa et al., *Psychological Support, Puberty Suppression, and Psychosocial Functioning in Adolescents with Gender Dysphoria*, 12 J. SEXUAL MED. 2206–2214 (2015), <https://pubmed.ncbi.nlm.nih.gov/26556015>; Annelou L.C. de Vries et al., *Puberty Suppression In Adolescents With Gender Identity Disorder: A Prospective Follow-Up Study*, 8 J. SEXUAL MED. 2276–2283 (2011), <https://pubmed.ncbi.nlm.nih.gov/20646177>; Annelou L.C. de Vries et al., *Young Adult Psychological Outcome After Puberty Suppression and Gender Reassignment*, 134 PEDIATRICS 696–704 (2014), <https://pubmed.ncbi.nlm.nih.gov/25201798>; Laura E. Kuper, et al., *Body Dissatisfaction and Mental Health Outcomes of Youth on* (continued...)

therapy to treat adolescents with gender dysphoria.⁵⁸ These studies find positive mental health outcomes for those adolescents who received puberty blockers or hormone therapy, including statistically significant reductions in anxiety,

Gender-Affirming Hormone Therapy, 145 PEDIATRICS e20193006 (2020), <https://perma.cc/2HAT-GGFV>; Jack L. Turban et al., *Pubertal Suppression For Transgender Youth and Risk of Suicidal Ideation*, 145 PEDIATRICS e20191725 (2020), <https://perma.cc/B2UZ-YR3Q>; Anna I.R. van der Miesen, *Psychological Functioning in Transgender Adolescents Before and After Gender-Affirmative Care Compared with Cisgender General Population Peers*, 66 J. ADOLESCENT HEALTH 699–704 (2020); Diana M. Tordoff et al., *Mental Health Outcomes in Transgender and Nonbinary Youths Receiving Gender-Affirming Care*, 5 JAMA NETWORK OPEN e220978 (2022), <https://perma.cc/SBF4-B4D4>.

⁵⁸ See, e.g., Achille, *supra* note 57, at 1–5; Luke R. Allen et al., *Well-Being and Suicidality Among Transgender Youth After Gender-Affirming Hormones*, 7 CLINICAL PRAC. PEDIATRIC PSYCH. 302 (2019), <https://psycnet.apa.org/record/2019-52280-009>; Diane Chen et al., *Psychosocial Functioning in Transgender Youth after 2 Years of Hormones*, 388 NEW ENG. J. MED. 240-50 (2023), <https://www.nejm.org/doi/10.1056/NEJMoa2206297>; Diego Lopez de Lara et al., *Psychosocial Assessment in Transgender Adolescents*, 93 ANALES DE PEDIATRIA 41–48 (English ed. 2020), <https://perma.cc/AQ4G-YJ85>; de Vries, *Young Adult Psychological Outcome After Puberty Suppression and Gender Reassignment*, *supra* note 57; Rittakerttu Kaltiala et al., *Adolescent Development and Psychosocial Functioning After Starting Cross-Sex Hormones for Gender Dysphoria*, 74 NORDIC J. PSYCHIATRY 213 (2020), <https://doi.org/10.1080/08039488.2019.1691260>; Kuper, *supra* note 57; Amy E. Green et al., *Association of Gender-Affirming Hormone Therapy with Depression, Thoughts of Suicide, and Attempted Suicide Among Transgender and Nonbinary Youth*, J. ADOLESCENT HEALTH (2021), <https://doi.org/10.1016/j.jadohealth.2021.10.036>; Jack L. Turban et al., *Access to Gender-Affirming Hormones During Adolescence and Mental Health Outcomes Among Transgender Adults*, J. PLOS ONE (2022), <https://doi.org/10.1371/journal.pone.0261039>.

depression, and suicidal ideation.⁵⁹

For example, a 2020 study analyzed survey data from 89 transgender adults who had access to puberty blockers while adolescents and from more than 3,400 transgender adults who did not.⁶⁰ The study found that those who received puberty blocking treatment had lower odds of lifetime suicidal ideation than those who wanted puberty blocking treatment but did not receive it, even after adjusting for demographic variables and level of family support.⁶¹ Approximately *nine in ten* transgender adults who wanted puberty blocking treatment but did not receive it reported lifetime suicidal ideation.⁶² Additionally, a longitudinal study of nearly 50 transgender adolescents found that suicidality was decreased by a statistically significant degree after receiving gender-affirming hormone treatment.⁶³ A study published in January 2023, following 315 participants age 12 to 20 who received gender-affirming hormone treatment, found that the treatment was associated with

⁵⁹ The data likewise indicates that adults who receive gender-affirming care experience positive mental health outcomes. *See, e.g.,* Zoe Aldridge et al., *Long Term Effect of Gender Affirming Hormone Treatment on Depression and Anxiety Symptoms in Transgender People: A Prospective Cohort Study*, 9 *ANDROLOGY* 1808–1816 (2021), <https://perma.cc/543U-HL5P>.

⁶⁰ *See* Turban, *supra* note 57.

⁶¹ *See id.*

⁶² *See id.*

⁶³ *See* Allen, *supra* note 58.

decreased symptoms of depression and anxiety.⁶⁴

As another example, a prospective two-year follow-up study of adolescents with gender dysphoria published in 2011 found that treatment with puberty blockers was associated with decreased depression and improved overall functioning.⁶⁵ A six-year follow-up study of 55 individuals from the 2011 study found that subsequent treatment with hormone therapy followed by surgery in adulthood was associated with a statistically significant decrease in depression and anxiety.⁶⁶ “Remarkably, this study demonstrated that these transgender adolescents and young adults had a sense of well-being that was equivalent or superior to that seen in age-matched controls from the general population.”⁶⁷

As scientists and researchers, *amici* always welcome more research, including on this crucial topic. However, contrary to the assertions by the Legislature that these treatments are “harmful” or “experimental,”⁶⁸ the available data indicate that the gender-affirming medical care prohibited by the Healthcare Ban is effective for

⁶⁴ See Chen et al., *supra* note 58.

⁶⁵ See de Vries, *Puberty Suppression in Adolescents with Gender Identity Disorder: A Prospective Follow-Up Study*, *supra* note 57.

⁶⁶ de Vries, *Young Adult Psychological Outcome After Puberty Suppression and Gender Reassignment*, *supra* note 57.

⁶⁷ Rosenthal, *supra* note 18, at 586.

⁶⁸ S.B. 99 § 2, 68th Legis. (2023).

the treatment of gender dysphoria.

III. The Healthcare Ban Would Irreparably Harm Many Adolescents with Gender Dysphoria by Denying Them the Treatment They Need.

If enforced, the Healthcare Ban would prevent adolescents with gender dysphoria in Montana from accessing medical care that is designed to improve health outcomes and alleviate suffering, and that is grounded in science and endorsed by the medical community. As discussed in Part II.C, research shows that adolescents with gender dysphoria who receive puberty blockers and/or hormone therapy experience less depression, anxiety, and suicidal ideation. Several studies have found that hormone therapy is associated with reductions in the rate of suicide attempts and significant improvement in quality of life.⁶⁹ In light of this evidence supporting the connection between lack of access to gender-affirming medical care and lifetime suicide risk, banning such care can put patients' lives at risk.

CONCLUSION

For the foregoing reasons, this Court should affirm the District Court's

⁶⁹ See M. Hassan Murad et al., *Hormonal Therapy and Sex Reassignment: A Systematic Review and Meta-Analysis of Quality of Life and Psychosocial Outcomes*, 72 *Clinical Endocrinology* 214 (2010), <https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2265.2009.03625.x>; see also Chen et al., *supra* note 58; Turban et al., *supra* note 58.

preliminary injunction.

Dated: April 11, 2024

Respectfully submitted,

/s/ Jon M. Moyers

Jon Moyers
MOYERS KOHN LAW
3936 Ave. B, Suite D
Billings, MT 59102
Phone: (406) 655-4900
jon@jmoyerslaw.com

D. Jean Veta*
William R. Isasi*
COVINGTON & BURLING, LLP
One CityCenter
850 Tenth St., N.W.
Washington, D.C. 20001
Phone: (202) 662-6000
jveta@cov.com
wisasi@cov.com

Counsel for Amici Curiae
**Pro hac vice applications*
forthcoming

CERTIFICATE OF SERVICE

I hereby certify that foregoing Brief was served by electronic service upon counsel of record on April 11, 2024.

/s/ Jon Moyers

Jon Moyers
MOYERS KOHN LAW
3936 Ave. B, Suite D
Billings, MT 59102
Phone: (406) 655-4900
jon@jmoyerslaw.com

Local Counsel for Amici Curiae

CERTIFICATE OF COMPLIANCE

This Certificate of Compliance is made pursuant to Montana Rule of Appellate Procedure 11. I hereby certify that the foregoing Brief is proportionally spaced in size 14 Times New Roman font. Further, the foregoing Brief contains 4,770 words not including the table of contents, table of citations, Certificate of Service, or this Certificate of Compliance.

/s/ Jon Moyers

Jon Moyers
MOYERS KOHN LAW
3936 Ave. B, Suite D
Billings, MT 59102
Phone: (406) 655-4900
jon@jmoyerslaw.com

Local Counsel for Amici Curiae